

Type II Anyon Superconductivity

Jewan Kim and Jungdae Kim

Department of Physics, Seoul National University, Seoul, 151-742, KOREA

Taejin Lee

Department of Physics, Kangwon National University, Chuncheon 200-701, KOREA

Abstract

Assuming that the superconductivity which is described by the low energy effective action of the anyon system may be type II, we discuss its characteristics. We also study physical properties of the Chern-Simons vortices, which may be formed as the external magnetic field is applied to the system, such as their statistics, the free energy of single vortex and the interaction energy between two vortices.

PACS numbers: 74.20.-z, 74.65.+n, 11.15.Kc

In a seminal paper [1] Fetter, Hanna and Laughlin made an important step in the anyon superconductivity [2], showing that a neutral gas of anyons has a massless Goldstone mode and exhibits the Meissner effect. Their work has been further extended by Chen, Halperin, Wilczek and Witten [3]. Banks and Lykken [4] then subsequently found that it is possible to describe the system in terms of a standard Landau-Ginzburg order parameter if the radiatively induced Chern-Simons term cancels its bare counterpart. However, there are many subjects to be clarified before the anyon superconductivity is accepted as a realistic model for the high- T_c superconductivity. One of them is the issue raised by the recent observation [5] that all known high- T_c superconductors are of type II: The issue is whether the anyon superconductivity can be extended to describe the type II superconductivity. The purpose of this paper is to discuss the anyon superconductivity in the framework of the type II superconductivity theory and to study physical properties of the Chern-Simons vortices which play an important role in the type II anyon superconductivity.

In the Abelian-Higgs model with the Chern-Simons term [6] there exist vortex solutions of finite energy as in the case where the Chern-Simons term is absent. However, these Chern-Simons vortices have distinctive characters: Carrying both electric charge and magnetic flux, they acquire fractional spins and satisfy exotic statistics. The Chern-Simons term [7] introduces the Aharonov-Bohm [8] interaction between the vortices, endowing them with the magnetic fluxes. In this respect, the effect of the Chern-Simons term in the 2+1 dimensional Abelian-Higgs model is similar to that of the CP violating term $\theta F^* F$ in the four dimensional non-Abelian-Higgs model which leads to the dyon [9]. The Chern-Simons vortex persists [10,11] even in the Chern-Simons-Higgs model [12] where the Maxwell term is suppressed. Moreover, in such a case the vortex solutions can be self-dual, satisfying the Bogomol'nyi conditions [13], provided that the Higgs potential is of a certain form.

The anyon superconductor theory may be described by the Lagrangian density [4]

$$L = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \Pi f_{\mu\nu}f^{\mu\nu} + \kappa\epsilon^{\mu\nu\lambda}A_\mu(F_{\nu\lambda} + 2f_{\nu\lambda}) \quad (1)$$

where Π and κ are constants and a_μ is the statistical gauge field, $f_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu$. Its

non-relativistic version has been discussed in refs. [1,3]. The Lagrangian density may be derived as a low-energy effective action for the system of anyons with the electromagnetic interaction

$$L = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \kappa'\epsilon^{\mu\nu\lambda}a_\mu f_{\nu\lambda} + \bar{\psi}(D \cdot \gamma + m)\psi \quad (2)$$

where $D_\mu = \partial_\mu + ie(A_\mu - a_\mu)$. The relativistic effective Lagrangian for the fermion field ψ may result from some microscopic model such as that of ref. [14]. Integrating out the fermion field ψ results in the effective action

$$-i \ln \text{Det}(D \cdot \gamma + m), \quad (3)$$

which has the low-energy expansion [7,15] given by

$$-\frac{1}{48\pi} \frac{e^2}{|m|} (F_{\mu\nu} - f_{\mu\nu})(F^{\mu\nu} - f^{\mu\nu}) + \frac{e^2}{16\pi} \frac{m}{|m|} \epsilon^{\mu\nu\lambda} (A_\mu - a_\mu)(F_{\nu\lambda} - f_{\nu\lambda}) + \dots \quad (4)$$

We may remove $F^{\mu\nu}F_{\mu\nu}$ and $f^{\mu\nu}F_{\mu\nu}$ terms in the above low-energy expansion by shifting the statistical gauge field, $a_\mu \rightarrow a_\mu + A_\mu$. Thus, the low-energy effective action for the system of anyons with the electromagnetic interaction is written as $\int L_{\text{eff}}$ where

$$L_{\text{eff}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{48\pi} \frac{e^2}{|m|} f_{\mu\nu}f^{\mu\nu} + \frac{e^2}{16\pi} \frac{m}{|m|} \epsilon^{\mu\nu\lambda} a_\mu f_{\nu\lambda} + \kappa'\epsilon^{\mu\nu\lambda} (A_\mu + a_\mu)(F_{\nu\lambda} + f_{\nu\lambda}). \quad (5)$$

From the effective action Eq.(5) we may read the condition [4] for the massless Goldstone mode to exist: The radiatively induced Chern-Simons term for the statistical gauge field cancels its bare term

$$\kappa' = -\frac{e^2}{16\pi} \frac{m}{|m|}. \quad (6)$$

This cancellation, which would not occur if ψ is nonrelativistic, holds up exactly all higher order due to the non-renormalization theorem [16,17] for the Chern-Simons term. Note that the fermion ψ has a statistical phase $\theta = \pm 2\pi$ with the above value of κ' , so the Chern-Simons interaction does not alter the statistics of ψ . Comparison of Eq.(1) and Eq.(5) yields the two constants Π and κ

$$\Pi = \frac{1}{48\pi} \frac{e^2}{|m|}, \quad \kappa = -\frac{e^2}{16\pi} \frac{m}{|m|}. \quad (7)$$

The field equations of motion which result from the Lagrangian density Eq.(1) are

$$\partial_\nu F^{\mu\nu} - 2\kappa\epsilon^{\mu\nu\lambda} (F_{\nu\lambda} + f_{\nu\lambda}) = 0, \quad (8a)$$

$$\Pi\partial_\nu f^{\mu\nu} - \kappa\epsilon^{\mu\nu\lambda}\partial_\nu A_\lambda = 0. \quad (8b)$$

Eq.(8b) is satisfied if we take

$$f^{\mu\nu} = \frac{\kappa}{\Pi}\epsilon^{\mu\nu\lambda}A_\lambda + c\epsilon^{\mu\nu\lambda}\partial_\lambda\phi \quad (9)$$

where c is a non-zero constant and will be chosen $\frac{1}{2\Pi}$ for convenience. Then the equations of motion are rewritten as

$$\partial_\nu F^{\mu\nu} - 2\kappa\epsilon^{\mu\nu\lambda}F_{\nu\lambda} - \frac{2\kappa}{\Pi}(\partial^\mu\phi + 2\kappa A^\mu) = 0, \quad (10a)$$

$$\partial_\mu(\partial^\mu\phi + 2\kappa A^\mu) = 0. \quad (10b)$$

Eq.(10b) is the Bianchi identity

$$\partial_\mu^* f^\mu = \partial_\mu \left(\frac{1}{2}\epsilon^{\mu\nu\lambda}f_{\nu\lambda} \right) = 0. \quad (11)$$

One easily identifies Eq.(10) as the equations of motion which follow from the Lagrangian

$$L = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \kappa\epsilon^{\mu\nu\lambda}A_\mu F_{\nu\lambda} + \frac{1}{2\Pi}(\partial^\mu\phi + 2\kappa A^\mu)^2. \quad (12)$$

The obtained Lagrangian Eq.(12) is similar to that of the well-known Landau-Ginzburg model [18] or equivalently the Abelian-Higgs model. If it were not for the Chern-Simons term, it would be considered as the Abelian-Higgs model

$$L = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}|(\partial_\mu + i2eA_\mu)\chi|^2 + V(|\chi|) \quad (13)$$

in the London limit [19] where the expectation value of $|\chi|$ is kept constant. The London limit is particularly useful to describe the vortex of type II superconductor [20,21] in that the coherence length ξ characterizing the size of the core of the vortex is much smaller than

other characteristic lengths, and the variation of the order parameter $|\chi|$ is negligible outside of the core. Note that, however, the vortex solutions apparently look singular in the London limit. This observation suggests that, although the vortex solutions to be explored in what follows are singular, they may be relevant to the anyon superconductivity. Therefore, it is meaningful to study the vortex solution of the Lagrangian Eq.(12) and to discuss their roles in the anyon superconductivity in the context of the type II superconductor theory. Recalling that the most physical properties of the vortex in the type II superconductor can be well described without invoking the inside structure of the core, we expect that we get a fairly good description of the type II anyon superconductor.

For a static solution the equations of motion Eq.(10) reduce to

$$\partial_j F^{ij} - 4\kappa \epsilon^{ij} \partial_j A_0 - \frac{2\kappa}{\Pi} \left(\partial^i \phi + 2\kappa A^i \right) = 0 \quad (14a)$$

$$\nabla^2 A_0 - \frac{4\kappa^2}{\Pi} A_0 + 4\kappa H = 0 \quad (14b)$$

$$\partial_i \left(\partial^i \phi + 2\kappa A^i \right) = 0 \quad (14c)$$

where $H = -\frac{1}{2}\epsilon^{ij}F_{ij}$. Choosing the Coulomb gauge $\partial^i A_i = 0$, we set $A_i = \epsilon_{ij}\partial^j f$. Taking the Ansatz for the axially symmetric vortex solution of vorticity n

$$A_0 = A_0(r), \quad f = f(r), \quad \text{and} \quad \phi = \frac{\kappa}{e}n\theta, \quad (15)$$

we can further reduce the equations of motion to

$$\left(\nabla^2 - \mu^2 \right) f - 4\kappa A_0 = \left(\frac{n\mu^2}{2e} \right) \ln r \quad (16a)$$

$$\left(\nabla^2 - \mu^2 \right) A_0 - 4\kappa \nabla^2 f = 0 \quad (16b)$$

where $\mu^2 = \frac{4\kappa^2}{\Pi}$. The reason to take the third condition in the Ansatz Eq.(15) is that we may identify ϕ as a Goldstone mode, i.e.,

$$\chi = |\chi| e^{i\frac{e}{\kappa}\phi}, \quad (17)$$

which follows from the comparison of Eq.(12) with Eq.(13). Concurrently we find that

$$|\chi| = \frac{|\kappa|}{e\sqrt{\Pi}} = \sqrt{\frac{3|m|}{16\pi}}. \quad (18)$$

With some algebra we can solve exactly the differential equations (16)

$$f(r) = -\frac{n}{4e(\Pi+1)} \left[\frac{1}{1+\Delta} (K_0(a_+r) + \ln r) + \frac{1}{1-\Delta} (K_0(a_-r) + \ln r) \right] \quad (19a)$$

$$A_0(r) = -\frac{\kappa n \Delta}{2e\Pi} [K_0(a_+r) - K_0(a_-r)] \quad (19b)$$

where $a_{\pm} = 2|\kappa|(\Delta^{-1} \pm 1)$ and $\Delta = \sqrt{\frac{\Pi}{\Pi+1}} < 1$. The electric charge density associated with the vortex is simply $j_0 = \mu^2 A_0$. Here K_0 is the modified Bessel function which we represent as

$$K_0(\nu|\mathbf{x} - \mathbf{x}'|) = -\int \frac{d^2k}{2\pi} \frac{e^{-i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}')}}{k^2 + \nu^2}. \quad (20)$$

The total magnetic flux Φ and the electric charge Q of the vortex which the solution Eq.(19) describes are

$$\Phi = \int d^2x H = -2\pi r \frac{\partial f}{\partial r} \Big|_{r=\infty} = \frac{\pi n}{e}, \quad (21a)$$

$$Q = \int d^2x \mu^2 A_0 = -\frac{ne}{4} \frac{m}{|m|} = 4\kappa\Phi. \quad (21b)$$

The relation between Φ and Q can be read from the equations of motion, so it applies to general static solutions. According to Eq.(21) and Eq.(7) the vortex itself is an anyon with the statistical parameter

$$\theta_s = \frac{Q\Phi}{2} = -\frac{n^2}{8} \frac{m}{|m|} \pi. \quad (22)$$

Note that the statistical parameter for the vortex of vorticity n depends upon the charge of χ which we assign to it. If we choose the charge of χ to be pe , the statistical parameter for the vortex would read as

$$\theta_s = -\frac{n^2}{2p^2} \frac{m}{|m|} \pi.$$

We choose here $2e$ for the charge of χ , because we consider χ corresponds to the bound state of a pair of ψ .

Making use of the analytic exact expression Eq.(19), we can evaluate the electric and magnetic fields that the vortex carries

$$\mathbf{E}(\mathbf{x}) = \frac{\mathbf{x}}{r} \frac{\kappa^2 n}{2\Pi} [(1 + \Delta)K_1(a_+ r) - (1 - \Delta)K_1(a_- r)], \quad (23a)$$

$$\begin{aligned} H(\mathbf{x}) &= -\nabla^2 f(r) \\ &= \frac{\kappa^2 n}{e\Pi} [(1 + \Delta)K_0(a_+ r) + (1 - \Delta)K_0(a_- r)]. \end{aligned} \quad (23b)$$

By means of the asymptotic expressions of the modified Bessel function

$$K_0(x) = -\ln \frac{x}{2} - \gamma \quad \text{for } x \ll 1, \quad K_0(x) = \sqrt{\frac{\pi}{2x}} e^{-x} \quad \text{for } x \gg 1,$$

we can read the behaviors of the various physical quantities near the core or far from it. In particular, we find that the magnetic field behaves as

$$H(\mathbf{x}) = \frac{\kappa^2 n}{e\Pi} \sqrt{\frac{\pi}{2r}} \left[\frac{(1 + \Delta)}{\sqrt{a_+}} e^{-a_+ r} + \frac{(1 - \Delta)}{\sqrt{a_-}} e^{-a_- r} \right] \quad (24)$$

for $r \gg \delta_{\pm} > \xi$, so the decrease of the field is characterized by two penetration depth parameters $\delta_{\pm}$. The analytic expression Eq.(19) also enables us to evaluate the free energy (or the mass of the vortex) of a single vortex with vorticity n

$$\begin{aligned} F_n &= \int d^2x \left(\frac{1}{2} (H^2 + E^2) + \frac{1}{2\Pi} \left(4\kappa^2 A_0^2 + (\partial^i \phi + 2\kappa A^i)^2 \right) \right) \\ &= \frac{1}{2\pi} \left(\frac{en\Delta}{16\Pi} \right)^2 \left[\frac{1}{1 + \Delta} \ln(\delta_+/\xi) + \frac{1}{1 - \Delta} \ln(\delta_-/\xi) \right]. \end{aligned} \quad (25)$$

Here we assume that the contribution from the inside region of the core is negligible. (This is true if the inside of the core is described by Ginzburg-Landau type Lagrangian.) Note that the coherence length ξ cannot be determined by the Lagrangian Eq.(12) and may be fixed by some higher powers of fields which are suppressed in the Lagrangian Eq.(1). Here we take it as a phenomenological parameter characterizing the theory. However, since F_n has logarithmic accuracy, it is insensitive to the precise value of ξ . Equation (25) indicates that the mass F_n is proportional to n^2 as in the case of the usual type II superconductor vortex, therefore, formation of vortex lines with one flux quantum is the most favourable.

The interaction energy of two vortices at a distance $d \gg \xi$ can be calculated similarly. Thanks to the linearity of the equations of motion Eq.(14) which are valid outside of the core, the system of two vortices at a distance $d \gg \xi$ can be described by a superposition of the fields of individual vortices. To evaluate the interaction energy F_{12} of two vortices we calculate the total free energy of the system, then subtract the masses of the individual vortices from it. Using the equations of motion Eq.(14), we rewrite the free energy as

$$\begin{aligned} F(H, A_0) &= \frac{1}{2} \int d^2x \left(H^2 + E^2 + \mu^2 A_0^2 + \frac{1}{\mu^2} (\partial_i (H + 4\kappa A_0))^2 \right) \\ &= \frac{1}{2} \int_S d\mathbf{l} \cdot \frac{\nabla}{2} \left(A_0^2 + \frac{1}{\mu^2} (H + 4\kappa A_0)^2 \right) \end{aligned} \quad (26)$$

where S is the boundary of the two dimensional space. For the system under discussion the total free energy is given by $F(H^T, A_0^T)$ with $H^T(\mathbf{r}) = H(\mathbf{r} - \mathbf{r}_1) + H(\mathbf{r} - \mathbf{r}_2)$, $A_0^T(\mathbf{r}) = A_0(\mathbf{r} - \mathbf{r}_1) + A_0(\mathbf{r} - \mathbf{r}_2)$ and S composed of two small circles, say, S_1 and S_2 , enclosing the cores of the vortices. Here $A_0(\mathbf{r})$ and $H(\mathbf{r})$ are the fields describing a single vortex located at $\mathbf{r}$ of which explicit expressions are given by Eq.(19b) and Eq.(23b). Accordingly the interaction energy F_{12} is given by

$$\begin{aligned} F_{12}(d) &= \frac{1}{2} \int_{S_1} d\mathbf{l} \cdot \nabla \left(A_0 + \frac{1}{\mu^2} (H + 4\kappa A_0) \right) (\mathbf{r} - \mathbf{r}_1) \\ &\quad \left(A_0 + \frac{1}{\mu^2} (H + 4\kappa A_0) \right) (\mathbf{r} - \mathbf{r}_2) + (1 \leftrightarrow 2) \\ &= \frac{1}{\pi} \left(\frac{en}{16} \right)^2 \frac{(1 - \Delta^2)}{\Delta^2} [(1 - \Delta)K_0(a_+d) + (1 + \Delta)K_0(a_-d)]. \end{aligned} \quad (27)$$

The interaction is always repulsive as in the usual case of the type II superconductor.

A few brief comments are in order on the critical fields of the anyon superconductivity model. The clear distinction between type I and type II superconductors lies in their magnetic properties. A typical type II superconductor has two critical fields, H_{c1} and H_{c2} : When the applied magnetic field H is smaller than H_{c1} , the superconductor exhibits the perfect Meissner effect. As the applied field exceeds H_{c1} , the vortices begin to form and for $H_{c1} < H < H_{c2}$ the specimen enters a new state called mixed state where a lattice of supercurrent vortices is formed. Finally for $H > H_{c2}$ the specimen becomes normal. The

critical field H_{c1} can be determined by making use of the analytic expression of the free energy of a single vortex Eq.(25). If the free energy of the single F_n vortex is smaller than the magnetic energy F_M , the formation of a vortex is favourable. The vortex of the vorticity n first appears when

$$F_n = \Phi_n H_{c1}. \quad (28)$$

This yields H_{c1}

$$H_{c1} = \frac{eF_n}{\pi n} \simeq \frac{3en|m|}{32\pi} (\ln(\delta_+/\xi) + \ln(\delta_-/\xi)). \quad (29)$$

But the critical field H_{c2} cannot be determined by the given Lagrangian Eq.(1). In order to evaluate it we may need the terms of higher powers in fields which are suppressed in the effective action.

We conclude this paper with some remarks on what remains to be discussed further. In this paper we explored the effective action for the anyon superconductor in the context of the type II superconductivity theory and showed that it is certainly possible for the anyon superconductor to be type II. However, there are several subjects to be clarified in the future works. We expect that the magnetization curve of the anyon superconductor near the critical field H_{c1} may radically differ from that of regular type II superconductor. The reason is that there are two penetration depth parameters instead of one and the vortex of the lowest free energy satisfies the fractional statistics rather than the bose statistics. Our discussion of this paper shows that it is urgent to investigate the question of whether the terms of higher powers in fields may improve the theory (e.g. they render the vortex solutions regular) and help us to determine ξ and H_{c2} . Last but not least, it is also important to clearly understand the effect of the four-fermi interaction [22], in the anyon superconductor theory, which plays an essential role in the ordinary BCS type superconductor theory.

ACKNOWLEDGMENTS

This work was supported in part by non-directed research fund, Korea Research Foundation (1992) and in part by BSRI-92-206.

REFERENCES

- [1] A. Fetter, C. Hanna, and R. Laughlin, Phys. Rev. **B39** (1989) 9679.
- [2] For a review see, J. Lykken, J. Sonnenschein and N. Weiss, Int. J. Mod. Phys. **A6** (1991) 5155.
- [3] Y.-H. Chen, B. I. Halperin, F. Wilczek and E. Witten, Int. J. Mod. Phys. **B3** (1989) 1001.
- [4] T. Banks and J. D. Lykken, Nucl. Phys. **B336** (1990) 500.
- [5] G. Burns, *High-Temperature Superconductivity* (Academic Press, 1992).
- [6] H. J. de Vega and F. A. Schaposnik, Phys. Rev. Lett. **56** (1986) 2564; C. N. Kumar and A. Khare, Phys. Lett. **178B** (1986) 395.
- [7] S. Deser, R. Jackiw, and R. Templeton, Ann. Phys. **140** (1982) 327.
- [8] Y. Aharonov and D. Bohm, Phys. Rev. **115** (1959) 485.
- [9] B. Julia and A. Zee, Phys. Rev. **D11** (1975) 2227.
- [10] J. Hong, Y. Kim and P. Y. Pac, Phys. Rev. Lett. **64** (1990) 2230.
- [11] R. Jackiw and E. Weinberger, Phys. Rev. Lett. **64** (1990) 2234.
- [12] S. Deser and Z. Yang, Mod. Phys. Lett. **A4** (1989) 2123.
- [13] E. B. Bogomol'nyi, Sov. J. Nucl. Phys. **24** (1976) 449; H. deVega and F. Schaposnik, Phys. Rev. **D14** (1976) 1100; L. Jacobs and C. Rebbi, Phys. Rev. **B19** (1978) 4486.
- [14] F. D. M. Haldane, Phys. Rev. Lett. **61** (1988) 2015.
- [15] T. Lee, J. Math. Phys. **27** (1986) 2434; S. Deser and A. N. Redlich, Phys. Rev. Lett. **61** (1988) 1541; I. Kogan and G. W. Semenoff, Nucl. Phys. **B368** (1992) 718.
- [16] Y. Kao and M. Suzuki, Phys. Rev. **D31** (1985) 2137; M. Bernstein and T. Lee, Phys.

- Rev. **D32** (1985) 1020.
- [17] T. Lee, Phys. Lett. **171B** (1986) 247; S. Coleman and B. Hill, Phys. Lett. **159B** (1985) 184; G. W. Semenoff, P. Sodano, and Y.-S. Wu, Phys. Rev. Lett. **62** (1989) 715; D. K. Hong, T. Lee and S. H. Park, Phys. Rev. **D48** (1993) 3918.
- [18] V. L. Ginzburg and L. D. Landau, Zh. Eksp. Theor. Fiz. **20** (1950) 1064.
- [19] M. K. Prasad and C. M. Sommerfield, Phys. Rev. Lett. **35** (1975) 760.
- [20] A. A. Abrikosov, Sov. Phys. JETP **5** (1957) 1174.
- [21] E. M. Lifshitz and L. P. Pitaevskii, Statistical Physics (Part 2) (Pergamon Press, Oxford, 1980); D. Saint-James, G. Sarma and E. J. Thomas, Type II Superconductivity (Pergamon Press, Oxford, 1969).
- [22] B. Rosenstein, B. J. Warr and S. H. Park, Phys. Rev. Lett. **62** (1989) 1433; G. W. Semenoff, P. Sodano and Y. S. Wu, Phys. Rev. Lett. **62** (1989) 715; J. Chay, D. K. Hong, T. Lee and S. H. Park, Phys. Rev. **D48** (1993) 909.